

HOSTED BY

Available online at www.sciencedirect.com**ScienceDirect**journal homepage: www.elsevier.com/locate/ajps**Original Research Paper****IL 10****Doxorubicin: The dogma and the bone****Jia Y. Wong^a, Pornsak Sriamornsak^b, Crispin R. Dass^{a,*}**^a School of Pharmacy, Curtin University, Bentley 6102, Australia^b Department of Pharmaceutical Technology, and Pharmaceutical Biopolymer Group (PBiG), Faculty of Pharmacy, Silpakorn University, Nakhon Pathom 73000, Thailand**A R T I C L E I N F O****Article history:**

Available online 23 November 2015

Keywords:

Doxorubicin

Cancer

Tumour

Apoptosis

Cardiotoxicity

Doxorubicin is an anthracycline drug used as frontline chemotherapy of osteosarcoma among other types of cancers. While quite effective in providing a good control over the tumours in some patients, it can cause significant toxicity to organs such as the heart [1]. Attempts to ameliorate this toxicity with agents such as dexrazoxane have been successful, but not without complications [2] as will be discussed in this talk. Pigment epithelium-derived factor (PEDF), a potent anti-angiogenic and osteogenetic protein [3], was used to reduce this side-effect, together with an analysis of levels of adenosine

triphosphate (ATP) and reactive oxygen species (ROS). The effects of low dose doxorubicin in a clinically relevant dosing model will be discussed.

Acknowledgements

The author acknowledges the support of a Curtin Academic 50 scheme and various members of his lab for data generated.

* E-mail address: Crispin.Dass@curtin.edu.au.

Peer review under responsibility of Shenyang Pharmaceutical University.
<http://dx.doi.org/10.1016/j.ajps.2015.10.016>

1818-0876/© 2016 The Authors. Production and hosting by Elsevier B.V. on behalf of Shenyang Pharmaceutical University. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).




Fig. 1 PEDF reduces extent of doxorubicin-induced cellular vacuolation in cardiomyocytes.

REFERENCES

- [1] Tacar O, Sriamornsak P, Dass CR. Doxorubicin: an update on anticancer molecular action, toxicity and novel drug delivery systems. *J Pharm Pharmacol* 2013;65:157-170.
- [2] Bryant J, Picot J, Levitt G, et al. Cardioprotection against the toxic effects of anthracyclines given to children with cancer: a systematic review. *Health Technol Assess* 2007;11:1-84.
- [3] Alcantara MB, Nemazannikova N, Elahy M, et al. Pigment epithelium-derived factor upregulates collagen I and downregulates matrix metalloproteinase 2 in osteosarcoma cells, and colocalises to collagen I and heat shock protein 47 in fetal and adult bone. *J Pharm Pharmacol* 2014;66:1586-1592.